

Preparing for your treatment with TYVASO

You may have side effects...

Your healthcare team will help you understand your medication, including possible side effects. The most common side effects are cough, headache, throat irritation and pain, and dizziness. Having a plan to manage potential side effects can help you feel prepared and stay on track to get the full benefits of your medicine.

Be sure to contact your healthcare team about things you can do to help manage side effects while continuing treatment, such as:



COUGH

Review proper technique

Before treatment, eat a small amount of yogurt, honey, or peanut butter, or drink very cold or warm water



HEADACHE

Use an over-the-counter pain reliever, like acetaminophen



THROAT IRRITATION

If needed, use a throat spray to numb your throat before treatment

Patient Pro-Tip

“Everyone’s side effects may be different, so it’s important to contact your doctor. I will cough after each session, but it’s a minor cough. To manage the cough, I usually will take a drink of water.”

– Tracey, Living with PAH and Taking TYVASO DPI

Did you Know?

Most TYVASO side effects are temporary and may improve as your body gets used to treatment.

Talk to your healthcare provider to see if slowing the pace at which you increase your dose or adjusting other medications you’re taking could help.

These tips may help you manage common side effects with TYVASO, based on healthcare provider experience. They are not intended to be considered medical advice.



Review proper technique for your device in the Instructions for Use manuals in the pocket and/or by viewing the How to Use videos at [TYVASO.com](https://www.tyvaso.com).

Important Safety Information

Before you take TYVASO or TYVASO DPI, tell your healthcare provider about all of your medical conditions, including if you:

- Have low blood pressure
- Have or have had bleeding problems
- Have asthma or chronic obstructive pulmonary disease (COPD)

Please see Important Safety Information continued on reverse, and the accompanying Full Prescribing Information and Instructions for Use for TYVASO and TYVASO DPI in pocket.

TYVASO DPI
(treprostinil) INHALATION POWDER

TYVASO
(treprostinil) INHALATION SOLUTION

Important Safety Information (continued)

Before you take TYVASO or TYVASO DPI, tell your healthcare provider about all of your medical conditions, including if you: (continued)

- Are pregnant or plan to become pregnant. It is not known if either product will harm your unborn baby.
- Are breastfeeding or plan to breastfeed. It is not known if either product passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Either product and other medicines may affect each other.

Especially tell your healthcare provider if you take:

- Medicines used to treat high blood pressure or heart disease
- Medicines that decrease blood clotting (anticoagulants)
- Water pills (diuretics)
- Gemfibrozil (Lopid®) or rifampin (Rimactane®, Rifadin®, Rifamate®, Rifater®)

What are the possible side effects of TYVASO or TYVASO DPI?

Both products can cause **serious side effects**, including:

- **Low blood pressure** (symptomatic hypotension). If you have low blood pressure, either product may lower your blood pressure more.
- **Bleeding problems.** Either product may increase the risk of bleeding, especially in people who take blood thinners (anticoagulants).

The most common side effects of both products are cough, headache, throat irritation and pain, nausea, reddening of the face and neck (flushing), fainting or loss of consciousness, dizziness, diarrhea, and shortness of breath. Like other inhaled prostaglandins, you may have trouble breathing after taking TYVASO or TYVASO DPI because it may cause the muscles around your airway to tighten (bronchospasm). These are not all the possible side effects. Call your doctor for medical advice about side effects or if you have trouble breathing.

You may report side effects to the FDA at www.fda.gov/MedWatch or call 1-800-FDA-1088.

What are TYVASO and TYVASO DPI?

TYVASO (treprostinil) Inhalation Solution and TYVASO DPI (treprostinil) Inhalation Powder are prescription medicines used in adults to treat:

- Pulmonary arterial hypertension (PAH; WHO Group 1), which is high blood pressure in the arteries of your lungs. TYVASO or TYVASO DPI can improve the ability to exercise. Your ability to exercise decreases 4 hours after taking TYVASO or TYVASO DPI. It is not known if TYVASO or TYVASO DPI is safe and effective in children under 18 years of age.
- Pulmonary hypertension associated with interstitial lung disease (PH-ILD; WHO Group 3), which is high blood pressure in the lungs due to inflammation and sometimes scarring in the lungs. TYVASO or TYVASO DPI can improve the ability to exercise.

The risk information provided here is not comprehensive. To learn more about TYVASO or TYVASO DPI, talk with your healthcare provider. Please see Full Prescribing Information for TYVASO or TYVASO DPI, Instructions for Use manuals for TD-100 and TD-300 TYVASO® Inhalation System and TYVASO DPI® Inhalation Powder, and additional information at www.TYVASO.com or call 1-844-UNITHER (1-844-864-8437).

TYVISIconSEP2024